



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: October 21, 2015

ACCEPTED: November 29, 2015

PUBLISHED: December 15, 2015

Exact partition function in $U(2) \times U(2)$ ABJM theory deformed by mass and Fayet-Iliopoulos terms

Jorge G. Russo^{a,b} and Guillermo A. Silva^{c,d}

^a*Institució Catalana de Recerca i Estudis Avançats (ICREA),
Pg. Lluís Companys, 23, 08010 Barcelona, Spain*

^b*Department ECM, Institut de Ciències del Cosmos,
Universitat de Barcelona, Martí Franquès, 1, 08028 Barcelona, Spain*

^c*Instituto de Física La Plata, CONICET & Departamento de Física,
Universidad Nacional de La Plata, C.C. 67, 1900 La Plata, Argentina*

^d*Abdus Salam International Centre for Theoretical Physics, Associate Scheme,
Strada Costiera 11, 34151 Trieste, Italy*

E-mail: jorge.russo@icrea.cat, silva@fisica.unlp.edu.ar

ABSTRACT: We exactly compute the partition function for $U(2)_k \times U(2)_{-k}$ ABJM theory on S^3 deformed by mass m and Fayet-Iliopoulos parameter ζ . For $k = 1, 2$, the partition function has an infinite number of Lee-Yang zeros. For general k , in the decompactification limit the theory exhibits a quantum (first-order) phase transition at $m = 2\zeta$.

KEYWORDS: Matrix Models, Supersymmetric gauge theory, AdS-CFT Correspondence, Chern-Simons Theories

ARXIV EPRINT: [1510.02957](https://arxiv.org/abs/1510.02957)

Contents

1	Introduction	1
2	Residue integration	2
3	Summary of results and limits	5
4	The special case $m_2 = 0$	8

1 Introduction

The dynamics of two coincident M2 branes on the orbifold $\mathbb{R}^8/\mathbb{Z}_k$ is described by ABJM theory, three-dimensional $U(2)_k \times U(2)_{-k}$ supersymmetric Chern-Simons theory with bi-fundamental matter [1]. For this particular gauge group, the ABJM theory has $\mathcal{N} = 8$ superconformal symmetry and is in fact equivalent to Gustavsson-Bagger-Lambert theory [2, 3]. The partition function for the theory on \mathbb{S}^3 can be computed by supersymmetric localization [4, 5]. This theory can be deformed, preserving $\mathcal{N} = 4$ supersymmetry, by adding mass and Fayet-Iliopoulos (FI) parameters m, ζ , and the localization technique then reduces the full supersymmetric functional integral to the matrix integral [5]

$$Z = \frac{1}{4} \int \frac{d^2\mu}{(2\pi)^2} \frac{d^2\nu}{(2\pi)^2} \frac{\sinh^2 \frac{\mu_1 - \mu_2}{2} \sinh^2 \frac{\nu_1 - \nu_2}{2}}{\prod_{i,j} \cosh \left(\frac{\mu_i - \nu_j + m}{2} \right) \cosh \left(\frac{\mu_i - \nu_j - m}{2} \right)} e^{\frac{ik}{4\pi} \sum_i (\mu_i^2 - \nu_i^2) - \frac{ik}{2\pi} \zeta \sum_i (\mu_i + \nu_i)} \quad (1.1)$$

where $i, j = 1, 2$. The parameter ζ represents a Fayet-Iliopoulos parameter for the diagonal $U(1)$ subgroup, whereas m corresponds to a mass for the chiral multiplets. The partition function should be understood as a function $Z(2\zeta, m; k)$, but for ease of presentation we will omit its arguments unless needed. For $k = 1$, the theory is mirror dual to $\mathcal{N} = 4$ supersymmetric super Yang-Mills theory with gauge group $U(2)$ coupled to a single fundamental hypermultiplet and a single adjoint hypermultiplet [5].

By shifting the integration variables, $x \equiv \mu - \zeta$, $y \equiv \nu + \zeta$, the partition function becomes

$$Z = \frac{1}{4} \int \frac{d^2x}{(2\pi)^2} \frac{d^2y}{(2\pi)^2} \frac{\sinh^2 \frac{x_1 - x_2}{2} \sinh^2 \frac{y_1 - y_2}{2}}{\prod_{i,j} \cosh \frac{x_i - y_j + m_1}{2} \cosh \frac{x_i - y_j - m_2}{2}} e^{\frac{ik}{4\pi} \sum_i (x_i^2 - y_i^2)}, \quad (1.2)$$

where m_1, m_2 are

$$m_1 = m + 2\zeta \quad \text{and} \quad m_2 = m - 2\zeta. \quad (1.3)$$

Note that ζ has dimension of mass. We are using units where the radius R of the three-sphere is $R = 1$.

The purpose of this note is to explicitly carry out the integration in (1.2). In the $m = \zeta = 0$ case, the integral was computed in [6] (a discussion of the partition function in the more general ABJ case can be found in [7]). On the other hand, the m, ζ -deformed ABJM theory was studied in [8] using the Fermi-gas formulation [9] and at large N for the $U(N)_k \times U(N)_{-k}$ gauge group in [10] (with $\zeta = 0$) and in [11] (with general $m, \zeta \neq 0$), where phase transitions in the complex parameter space generated by m_1, m_2 and $g = 2\pi i/k$ were investigated. Our explicit formula will uncover some interesting physical properties of the mass-deformed system with gauge group $U(2)_k \times U(2)_{-k}$.

The partition function (1.2) manifests the $m_1 \leftrightarrow m_2$ symmetry or, equivalently, $\zeta \rightarrow -\zeta$. A less obvious symmetry is $m_2 \rightarrow -m_2$, or [8, 11]

$$Z(2\zeta, m; k) = Z(m, 2\zeta; k). \quad (1.4)$$

For the $k = 1$ case, this symmetry already appeared in [5], where it was also explained by the fact that the corresponding brane configuration is self-mirror. The symmetry implies, in particular, that a FI-deformation ζ on the massless theory is equivalent to a mass-deformation $m = 2\zeta$ in the theory with vanishing FI-parameter. The case $m = 2\zeta$ — representing a fixed point of this symmetry — is special, as we shall shortly see. In the dual $\mathcal{N} = 4$ supersymmetric super Yang-Mills theory, $m_2 = 0$ corresponds to coupling the theory to a massless adjoint hypermultiplet.

2 Residue integration

The partition function for the m, ζ -deformed ABJM theory with $U(N)_k \times U(N)_{-k}$ gauge group can be written in the following form [5, 11]

$$Z(2\zeta, m; k) = \sum_{\rho} (-1)^{\rho} \frac{1}{N!} \int d^N \tau \frac{e^{-ikm_2 \sum_i \tau_i}}{\prod_i \cosh(k\pi\tau_i) \cosh\left(\pi(\tau_i - \tau_{\rho(i)}) - \frac{m_1}{2}\right)}, \quad (2.1)$$

where the sum goes over permutations. The derivation uses a trigonometric identity, Fourier integrations and only holds for opposite Chern-Simons levels (see section 2 in [11] for details). For $N = 2$, the formula (2.1) then leads to the following expression

$$Z = \frac{1}{2}(Z_1 - Z_2), \quad (2.2)$$

with

$$Z_1 = \int d\tau_1 d\tau_2 \frac{e^{-ikm_2(\tau_1 + \tau_2)}}{\cosh(\pi k\tau_1) \cosh(\pi k\tau_2) \cosh^2\left(\frac{m_1}{2}\right)}, \quad (2.3)$$

and

$$Z_2 = \int d\tau_1 d\tau_2 \frac{e^{-ikm_2(\tau_1 + \tau_2)}}{\cosh(\pi k\tau_1) \cosh(\pi k\tau_2) \cosh\left(\pi(\tau_1 - \tau_2) - \frac{m_1}{2}\right) \cosh\left(\pi(\tau_1 - \tau_2) + \frac{m_1}{2}\right)}, \quad (2.4)$$

Using the identity

$$\frac{1}{\cosh^2 \frac{m_1}{2}} - \frac{1}{\cosh\left(\pi\tau - \frac{m_1}{2}\right) \cosh\left(\pi\tau + \frac{m_1}{2}\right)} = \frac{\operatorname{sech}^2 \frac{m_1}{2} \sinh^2 \pi\tau}{\cosh\left(\pi\tau - \frac{m_1}{2}\right) \cosh\left(\pi\tau + \frac{m_1}{2}\right)} \quad (2.5)$$

and the formula for the Fourier transform [11]

$$\int du \frac{e^{-ikm_2u}}{\cosh\left(\frac{\pi k}{2}(u+v)\right)\cosh\left(\frac{\pi k}{2}(u-v)\right)} = \frac{4\sin(km_2v)}{k\sinh(\pi kv)\sinh m_2}, \quad (2.6)$$

we obtain

$$Z = \frac{1}{k^2 \sinh(m_2) \cosh^2 \frac{m_1}{2}} \int du \frac{\sin(m_2u) \sinh^2 \frac{\pi u}{k}}{\sinh(\pi u) \cosh\left(\frac{\pi u}{k} - \frac{m_1}{2}\right) \cosh\left(\frac{\pi u}{k} + \frac{m_1}{2}\right)}. \quad (2.7)$$

In the limit $m_2 \rightarrow 0$, the partition function becomes

$$Z|_{m_2=0} = \frac{1}{k^2 \cosh^2 \frac{m_1}{2}} \int du \frac{u \sinh^2 \frac{\pi u}{k}}{\sinh(\pi u) \cosh\left(\frac{\pi u}{k} - \frac{m_1}{2}\right) \cosh\left(\frac{\pi u}{k} + \frac{m_1}{2}\right)}. \quad (2.8)$$

In the following, we compute the integrals (2.7), (2.8) by residue integration.

To compute (2.7) we follow the ideas in [6], where the partition function was computed in the case $m = \zeta = 0$.

Thus we start by writing the integrand as the product of two even functions f, g

$$Z = \frac{1}{k^2 \sinh(m_2) \cosh^2 \frac{m_1}{2}} \int du f(u)g(u), \quad (2.9)$$

with

$$f(u) = \frac{\sin m_2u}{\sinh \pi u}, \quad g(u) = \frac{\sinh^2 \frac{\pi u}{k}}{\cosh\left(\frac{\pi u}{k} - \frac{m_1}{2}\right) \cosh\left(\frac{\pi u}{k} + \frac{m_1}{2}\right)}. \quad (2.10)$$

Under the shift $u \rightarrow u + ik$ these functions transform as

$$\begin{aligned} f(u) &\rightarrow (-)^k \cosh(m_2k) f(u) + \text{odd function}, \\ g(u) &\rightarrow g(u) \end{aligned} \quad (2.11)$$

These properties imply that the integral in (2.9) along the curve $u = x + ik$ with $x \in \mathbb{R}$ will differ from the integration along the real axis by the factor $(-)^k \cosh(m_2k)$. Therefore, the rectangular contour composed by the real axis, two vertical segments and the displaced real axis $u = x + ik$ becomes appropriate for residue computation in the case $m_2 \neq 0$ (see figure 1).¹

The residues encircled by the contour comprise the ones arising from the poles of $f(z)$ located at $z = in$ with $n = 1, \dots, k$ and those of $g(z)$ located at $z_{\pm} = \pm \frac{m_1k}{2\pi} + i\frac{k}{2}$. The pole located at $z = ik$ does not contribute due to a double zero in the numerator of $g(z)$. Calling C the closed rectangular contour described above and $\mathcal{F}(z) = f(z)g(z)$ one finds

$$\begin{aligned} \oint_C dz \mathcal{F}(z) &= (1 - (-)^k \cosh(m_2k)) \int du \mathcal{F}(u) \\ &= 2\pi i \left[\sum_{n=1}^{k-1} \text{Res}_{z=in} \mathcal{F}(z) + \text{Res}_{z=z_{\pm}} \mathcal{F}(z) \right] \end{aligned}$$

¹It is easily seen that the vertical contours do not contribute when we push them to infinity.




Figure 1. Rectangular contour for residue computation. The poles on the imaginary axis $z = in$ with $n = 1, \dots, k-1$ arise from the f function, while those at $z_{\pm} = \pm \frac{m_1 k}{2\pi} + i \frac{k}{2}$ follow from the g function.

which gives

$$\int du \mathcal{F}(u) = \frac{2\pi i}{1 - (-)^k \cosh(m_2 k)} \left[-\frac{i}{\pi} \sum_{n=1}^{k-1} (-)^n \frac{\sin^2\left(\frac{n\pi}{k}\right) \sinh(m_2 n)}{\cosh\left(\frac{m_1}{2} - \frac{in\pi}{k}\right) \cosh\left(\frac{m_1}{2} + \frac{in\pi}{k}\right)} + R_k \right] \quad (2.12)$$

where

$$R_k = \begin{cases} (-)^{\frac{k}{2}} \frac{ik}{\pi} \frac{\coth \frac{m_1}{2} \sinh \frac{km_2}{2}}{\sinh \frac{km_1}{2}} \cos \frac{km_1 m_2}{2\pi}, & k \text{ even} \\ (-)^{\frac{k+1}{2}} \frac{ik}{\pi} \frac{\coth \frac{m_1}{2} \cosh \frac{km_2}{2}}{\cosh \frac{km_1}{2}} \sin \frac{km_1 m_2}{2\pi}, & k \text{ odd} \end{cases} \quad (2.13)$$

Case $m_2 = 0$, k odd. It is evident from (2.12) that the $m_2 \rightarrow 0$ limit of (2.9) is smooth, the result is

$$Z|_{m_2=0} = \frac{1}{k^2 \cosh^2 m} \left[\sum_{n=1}^{k-1} (-)^n \frac{n \sin^2\left(\frac{n\pi}{k}\right)}{\cosh\left(m - \frac{in\pi}{k}\right) \cosh\left(m + \frac{in\pi}{k}\right)} - (-)^{\frac{k+1}{2}} \frac{k^2 m \coth m}{\pi \cosh km} \right], \quad k \text{ odd} \quad (2.14)$$

where we have used $m_1 = 2m$.

Case $m_2 = 0$, k even. The factor multiplying the bracket in (2.12) prevents taking $m_2 \rightarrow 0$ in the even k case. To compute the integral in (2.8) we consider

$$I = \int du \tilde{f}(u) g(u), \quad (2.15)$$

with $g(u)$ as in (2.10) and

$$\tilde{f}(u) = \frac{i}{k} \frac{(u - ik/2)^2}{\sinh \pi u}.$$

Upon integration, the odd piece in \tilde{f} vanishes against $g(u)$ and therefore the partition function (2.8) can be written as

$$Z|_{m_2=0} = \frac{1}{k^2 \cosh^2 m} I \quad (2.16)$$

The shift $u \rightarrow u + ik$ in $\tilde{f}(u)$ gives

$$\tilde{f}(u) \rightarrow (-)^{k+1} \tilde{f}(-u).$$

As discussed below (2.11), this property makes the rectangular contour in figure 1 appropriate for computing I by residues.

For the residues analysis we should now consider the pole in $\tilde{f}(z)$ at the origin $z = 0$ but a zero in $g(z)$ eliminates it; along the same lines the residue from $z = ik/2$ is absent since a zero appears for \tilde{f} . Calling $\tilde{\mathcal{F}}(z) = \tilde{f}(z)g(z)$ one finds

$$\oint_C dz \tilde{\mathcal{F}}(z) = 2I,$$

on the other hand

$$\begin{aligned} \oint_C dz \tilde{\mathcal{F}}(z) &= 2\pi i \left[\sum_{n=0}^{k-1} \text{Res}_{z=in} \tilde{\mathcal{F}}(z) + \text{Res}_{z=z_{\pm}} \tilde{\mathcal{F}}(z) \right] \\ &= 2\pi i \left[\frac{i}{k\pi} \sum_{n=1}^{k-1} (-)^n \left(\frac{k}{2} - n \right)^2 \frac{\sin^2 \left(\frac{n\pi}{k} \right)}{\cosh \left(m - \frac{in\pi}{k} \right) \cosh \left(m + \frac{in\pi}{k} \right)} + \tilde{\mathcal{R}}_k \right]. \end{aligned} \quad (2.17)$$

where

$$\tilde{\mathcal{R}}_k = (-)^{\frac{k}{2}} \frac{2i(mk)^2 \coth(m) \sinh mk}{\pi^3 \cosh(2mk) - 1}$$

The $n = \frac{k}{2}$ term in the sum vanishes as expected. The final result is

$$\begin{aligned} Z|_{m_2=0} &= -\frac{1}{k \cosh^2 m} \cdot \\ &\quad \left[\sum_{n=1}^{k-1} (-)^n \left(\frac{n}{k} - \frac{1}{2} \right)^2 \frac{\sin^2 \left(\frac{n\pi}{k} \right)}{\cosh \left(m - \frac{in\pi}{k} \right) \cosh \left(m + \frac{in\pi}{k} \right)} + (-)^{\frac{k}{2}} \frac{2m^2 k \coth(m) \sinh mk}{\pi^2 \cosh(2mk) - 1} \right] \end{aligned} \quad (2.18)$$

3 Summary of results and limits

Thus we have obtained

$$Z = \frac{2}{k^2 \sinh(m_2)} \frac{1}{1 - (-1)^k \cosh(m_2 k)} (J_1 - J_2) \quad (3.1)$$

where

$$J_1 = \frac{1}{\cosh^2 \left(\frac{m_1}{2} \right)} \sum_{n=1}^{k-1} (-1)^n \frac{\sin^2 \left(\frac{n\pi}{k} \right) \sinh(m_2 n)}{\cosh \left(\frac{m_1}{2} - \frac{in\pi}{k} \right) \cosh \left(\frac{m_1}{2} + \frac{in\pi}{k} \right)} \quad (3.2)$$

and

$$J_2 = \begin{cases} (-)^{\frac{k}{2}} \frac{2k \sinh \frac{km_2}{2}}{\sinh(m_1) \sinh \frac{km_1}{2}} \cos \frac{km_1 m_2}{2\pi}, & k \text{ even} \\ (-)^{\frac{k+1}{2}} \frac{2k \cosh \frac{km_2}{2}}{\sinh(m_1) \cosh \frac{km_1}{2}} \sin \frac{km_1 m_2}{2\pi}, & k \text{ odd} \end{cases} \quad (3.3)$$

Using

$$\frac{2}{1 + \cosh \alpha} = \frac{1}{\cosh^2 \left(\frac{\alpha}{2} \right)}, \quad \frac{2}{1 - \cosh \alpha} = -\frac{1}{\sinh^2 \left(\frac{\alpha}{2} \right)}, \quad (3.4)$$

we can finally put the partition function in the form

$$Z|_{k \text{ even}} = -\frac{1}{k^2 \sinh(m_2) \sinh^2 \left(\frac{km_2}{2} \right)} (J_1 - J_2) \quad (3.5)$$

$$Z|_{k \text{ odd}} = \frac{1}{k^2 \sinh(m_2) \cosh^2 \left(\frac{km_2}{2} \right)} (J_1 - J_2) \quad (3.6)$$

In the formulas (3.5)–(3.6), the symmetry $m_1 \leftrightarrow m_2$ — which is manifest in the integral form (1.2) — is hidden. Interestingly, this symmetry is only recovered upon summation over n . On the other hand, the symmetry $m_2 \rightarrow -m_2$ is manifest.

Note that Z is real. While this is expected in a unitary theory, it is not generally the case in Chern-Simons theories (for a discussion, see [12]). In the present case, it is related to the fact the theory is a combination of two Chern-Simons theory with opposite levels.²

Consider, as particular examples, the important cases $k = 1, 2$. The partition functions take the form

$$Z|_{k=1} = \frac{2}{\sinh(m_1) \sinh(m_2) \cosh \left(\frac{m_1}{2} \right) \cosh \left(\frac{m_2}{2} \right)} \sin \left(\frac{m_1 m_2}{2\pi} \right), \quad (3.7)$$

$$Z|_{k=2} = \frac{2}{\sinh^2(m_1) \sinh^2(m_2)} \sin^2 \left(\frac{m_1 m_2}{2\pi} \right). \quad (3.8)$$

Now the symmetry $m_1 \leftrightarrow m_2$ has become manifest.

Note that the partition functions for $k = 1, 2$ have zeros. Restoring the R dependence, the zeros are located at

$$m_1 m_2 R^2 = 2\pi^2 n, \quad n = \pm 1, \pm 2, \dots \quad (3.9)$$

They represent Lee-Yang zeros (see, for example, [13]). In the infinite volume, $R \rightarrow \infty$, the zeros condense in a certain line, and a phase transition should emerge. The fact that the partition function has zeros seems to be related to the fact that the coupling, $g = 2\pi i/k$, is imaginary for real k . Indeed, from the general expressions (3.2)–(3.3) we see that the arguments of the sine and cosine functions in (3.7), (3.8) contain a factor π/k . If the coupling g is (unphysically) continued to the real line by taking $k \rightarrow ik$, the partition function zeros would then lie on the imaginary g -axis, in accordance with the Lee-Yang theorem (see [11] for a related discussion).

For the undeformed ABJM theory, the $k = 1$ case is of special interest, since it is conjectured to describe the dynamics of two M2 branes in eleven-dimensional Minkowski

²We thank Miguel Tierz for comments on this point.

spacetime. An interesting question is what is the origin of these Lee-Yang singularities in the brane realization.

The partition function $Z(2\zeta, m; k)$ does not have any zeros for $k > 2$. For higher values of k , the partition function becomes more involved, below we quote explicitly the $k = 3$ and $k = 4$ cases

$$Z|_{k=3} = \frac{2}{3} \frac{2 - \sin\left(\frac{3m_1m_2}{2\pi}\right) \operatorname{csch}\left(\frac{m_1}{2}\right) \operatorname{csch}\left(\frac{m_2}{2}\right)}{(\cosh m_1 + \cosh 2m_1)(\cosh m_2 + \cosh 2m_2)} \quad (3.10)$$

$$Z|_{k=4} = \frac{1 - \operatorname{sech}(m_1) - \operatorname{sech}(m_2) + \cos\left(\frac{2m_1m_2}{\pi}\right) \operatorname{sech}(m_2) \operatorname{sech}(m_1)}{8 \sinh^2 m_1 \sinh^2 m_2} \quad (3.11)$$

Note that the symmetry under the exchange $m_1 \leftrightarrow m_2$ is manifest.

Asymptotic formulas. Let us consider the limit of a large sphere, $mR \gg 1$, at fixed k . Assuming $m_1 > 0$, $m_2 > 0$ and restoring the R dependence, we find

$$Z|_{k=1} \sim 32 e^{-\frac{3}{2}(m_1+m_2)R} \sin\left(\frac{m_1m_2R^2}{2\pi}\right), \quad (3.12)$$

$$Z|_{k=2} \sim 32 e^{-2(m_1+m_2)R} \sin^2\left(\frac{m_1m_2R^2}{2\pi}\right), \quad (3.13)$$

$$Z|_{k>2} \sim \frac{64}{k^2} e^{-2(m_1+m_2)R} \sin^2\left(\frac{\pi}{k}\right). \quad (3.14)$$

The general asymptotic formula with arbitrary sign for m_2 and $m_2 \neq 0$, is obtained by replacing m_2 by $|m_2|$.

The absolute value implies a discontinuity in the first derivative of $F = -\ln Z$. This indicates a first-order phase transition in the parameter m_2 at $m_2 = 0$, i.e., when the two mass scales $m, 2\zeta$ cross. Explicitly, at large R , we have

$$F = 2(|m_1| + |m_2|)R + O(1), \quad k > 1. \quad (3.15)$$

Hence

$$\left. \frac{d\Delta F}{dm_2} \right|_{m_2=0} = 4R, \quad \Delta F \equiv F_{m_2>0} - F_{m_2<0}. \quad (3.16)$$

For $k = 1$ the discontinuity in the first derivative of ΔF is equal to $3R$, as can be seen from (3.12).

For the general theory with gauge group $U(N)_k \times U(N)_{-k}$, large N phase transitions in the complex parameter $Ng = 2\pi i N/k$ were studied in [10, 11]. These phase transitions require taking infinite volume and, at the same time, a strong coupling limit with fixed kR — a limit that already appeared in the context of supersymmetric $U(N)$ Chern-Simons theory with massive fundamental matter in [14, 15]. It should be noted that such decompactification limit is different from the present (more physical) limit of large R at fixed k .

Another interesting aspect of (3.14) is that it is in a form suitable for a weak coupling expansion in powers of $1/k$:

$$Z|_{k>2} \sim -\frac{32}{k^2} e^{-2(m_1+m_2)R} \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} \left(\frac{2\pi}{k}\right)^{2n}. \quad (3.17)$$

The perturbative expansion has an infinite radius of convergence. However, the original theory on the three-sphere of *finite* radius R has an asymptotic perturbative expansion, with $2n!$ asymptotic behavior for the $1/k^{2n}$ term. This can be seen by using the integral form (2.7) and generalizing the study of [16, 17] on the resurgence properties of the perturbation series of ABJM theory. Now, expanding the integrand in (2.7), one finds a series with finite radius of convergence determined by the poles of $\text{sech}(\pi u/k \pm m_1/2)$ in the complex u -plane. The integral over u then adds an extra $(2n)!$, leading to an asymptotic (but Borel summable) perturbation series.

4 The special case $m_2 = 0$

The $m_2 = 0$ case is special and must be considered separately. In particular, it represents the critical point in the phase transitions that arise in the decompactification limit. In section 2 we have obtained the following formulas:

Odd k :

$$Z|_{m_2=0} = \frac{1}{k^2 \cosh^2 m} \sum_{n=1}^{k-1} (-)^n \frac{n \sin^2 \frac{\pi n}{k}}{\cosh\left(m + \frac{i\pi n}{k}\right) \cosh\left(m - \frac{i\pi n}{k}\right)} + \frac{(-)^{\frac{k-1}{2}} 2m}{\pi \cosh(km) \sinh(2m)}. \quad (4.1)$$

Even k :

$$\begin{aligned} Z|_{m_2=0} = & \frac{1}{k \cosh^2 m} \sum_{n=1}^{k-1} (-)^{n+1} \left(\frac{n}{k} - \frac{1}{2}\right)^2 \frac{\sin^2\left(\frac{n\pi}{k}\right)}{\cosh\left(m - \frac{i\pi n}{k}\right) \cosh\left(m + \frac{i\pi n}{k}\right)} \\ & + (-)^{\frac{k}{2}+1} \frac{4m^2}{\pi^2} \frac{\sinh mk}{\sinh(2m)(\cosh(2mk) - 1)} \end{aligned} \quad (4.2)$$

In particular,

$$\begin{aligned} Z|_{k=1} &= \frac{2m}{\pi \cosh(m) \sinh(2m)}, \\ Z|_{k=2} &= \frac{2m^2}{\pi^2 \sinh^2(2m)}. \end{aligned} \quad (4.3)$$

Note that the partition function does not have zeros in this case.

Asymptotic formulas $m_2 = 0$. Consider again the limit of a large sphere, $mR \gg 1$, at fixed k , but now with $m_2 = 0$. We find

$$Z|_{k=1} \sim \frac{8mR}{\pi} e^{-3mR}, \quad (4.4)$$

$$Z|_{k=2} \sim \frac{8}{\pi^2} m^2 R^2 e^{-4mR}, \quad (4.5)$$

$$Z|_{k>2} \sim \frac{4}{k^2} e^{-4mR} \tan^2 \frac{\pi}{k}. \quad (4.6)$$

Note that these formulas differ from the asymptotic formulas (3.12)–(3.14) given above for $Z(m_1, m_2)$ at $m_2 = 0$. This is expected, since the latter were obtained by assuming $|m_1 R|, |m_2 R| \rightarrow \infty$.

Unlike the $m_2 \neq 0$ case, the perturbation series for this flat-theory limit has now finite radius of convergence $|\pi/k| < \pi/2$, therefore perturbation series is convergent for all $k > 2$, where the formula applies. On the other hand, just like the general $m_2 \neq 0$ case, the theory on a finite-radius \mathbb{S}^3 has an asymptotic perturbation series with $2n!$ asymptotic behavior.

Finally, it would be interesting to study supersymmetric Wilson loops in the present mass/FI deformed theory, along the lines of [18].

Acknowledgments

We would like to thank Miguel Tierz for useful comments. GAS acknowledge support from ICTP and would like to thank Department ECM of Universitat de Barcelona for warm hospitality during the early stages of this work. We acknowledge financial support from projects FPA2013-46570, 2014-SGR-104, PIP0595/13 CONICET and X648 UNLP.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] O. Aharony, O. Bergman, D.L. Jafferis and J. Maldacena, $\mathcal{N} = 6$ superconformal Chern-Simons-matter theories, $M2$ -branes and their gravity duals, *JHEP* **10** (2008) 091 [[arXiv:0806.1218](https://arxiv.org/abs/0806.1218)] [[INSPIRE](#)].
- [2] A. Gustavsson, Algebraic structures on parallel $M2$ -branes, *Nucl. Phys. B* **811** (2009) 66 [[arXiv:0709.1260](https://arxiv.org/abs/0709.1260)] [[INSPIRE](#)].
- [3] J. Bagger and N. Lambert, Gauge symmetry and supersymmetry of multiple $M2$ -branes, *Phys. Rev. D* **77** (2008) 065008 [[arXiv:0711.0955](https://arxiv.org/abs/0711.0955)] [[INSPIRE](#)].
- [4] A. Kapustin, B. Willett and I. Yaakov, Exact results for Wilson loops in superconformal Chern-Simons theories with matter, *JHEP* **03** (2010) 089 [[arXiv:0909.4559](https://arxiv.org/abs/0909.4559)] [[INSPIRE](#)].
- [5] A. Kapustin, B. Willett and I. Yaakov, Nonperturbative tests of three-dimensional dualities, *JHEP* **10** (2010) 013 [[arXiv:1003.5694](https://arxiv.org/abs/1003.5694)] [[INSPIRE](#)].
- [6] K. Okuyama, A note on the partition function of ABJM theory on S^3 , *Prog. Theor. Phys.* **127** (2012) 229 [[arXiv:1110.3555](https://arxiv.org/abs/1110.3555)] [[INSPIRE](#)].
- [7] H. Awata, S. Hirano and M. Shigemori, The partition function of ABJ theory, *Prog. Theor. Exp. Phys.* **2013** (2013) 053B04 [[arXiv:1212.2966](https://arxiv.org/abs/1212.2966)] [[INSPIRE](#)].
- [8] N. Drukker and J. Felix, 3d mirror symmetry as a canonical transformation, *JHEP* **05** (2015) 004 [[arXiv:1501.02268](https://arxiv.org/abs/1501.02268)] [[INSPIRE](#)].
- [9] M. Mariño and P. Putrov, ABJM theory as a Fermi gas, *J. Stat. Mech.* (2012) P03001 [[arXiv:1110.4066](https://arxiv.org/abs/1110.4066)] [[INSPIRE](#)].

- [10] L. Anderson and K. Zarembo, *Quantum phase transitions in mass-deformed ABJM matrix model*, *JHEP* **09** (2014) 021 [[arXiv:1406.3366](#)] [[INSPIRE](#)].
- [11] L. Anderson and J.G. Russo, *ABJM theory with mass and FI deformations and quantum phase transitions*, *JHEP* **05** (2015) 064 [[arXiv:1502.06828](#)] [[INSPIRE](#)].
- [12] C. Closset, T.T. Dumitrescu, G. Festuccia, Z. Komargodski and N. Seiberg, *Contact terms, unitarity and F-maximization in three-dimensional superconformal theories*, *JHEP* **10** (2012) 053 [[arXiv:1205.4142](#)] [[INSPIRE](#)].
- [13] C. Itzykson and J.M. Drouffe, *Statistical field theory. Vol. 1: From Brownian motion to renormalization and lattice gauge theory*, Cambridge University Press, Cambridge U.K. (1989) [[INSPIRE](#)].
- [14] A. Barranco and J.G. Russo, *Large- N phase transitions in supersymmetric Chern-Simons theory with massive matter*, *JHEP* **03** (2014) 012 [[arXiv:1401.3672](#)] [[INSPIRE](#)].
- [15] J.G. Russo, G.A. Silva and M. Tierz, *Supersymmetric $U(N)$ Chern-Simons-matter theory and phase transitions*, *Commun. Math. Phys.* **338** (2015) 1411 [[arXiv:1407.4794](#)] [[INSPIRE](#)].
- [16] J.G. Russo, *A note on perturbation series in supersymmetric gauge theories*, *JHEP* **06** (2012) 038 [[arXiv:1203.5061](#)] [[INSPIRE](#)].
- [17] I. Aniceto, J.G. Russo and R. Schiappa, *Resurgent analysis of localizable observables in supersymmetric gauge theories*, *JHEP* **03** (2015) 172 [[arXiv:1410.5834](#)] [[INSPIRE](#)].
- [18] S. Hirano, K. Nii and M. Shigemori, *ABJ Wilson loops and Seiberg duality*, *Prog. Theor. Exp. Phys.* **2014** (2014) 113B04 [[arXiv:1406.4141](#)] [[INSPIRE](#)].